

International Plant Nutrition Colloquium

(University of California, Davis)

Year 2009

Paper 1051

Difference in Acquisition of Soil Organic Nitrogen between Pak-Choi and Tomato

Toshihiro Watanabe *

Takumi Enomoto [†]

Miwa Okamoto [‡]

Michihiko Sakurai **

Takuro Shinano ^{††}

Mitsuru Osaki ^{‡‡}

*Hokkaido University

[†]Hokkaido University

[‡]National Agricultural Research Center for Hokkaido Region

**Hokkaido Central Agricultural Experiment Station

^{††}National Agricultural Research Center for Hokkaido Region

^{‡‡}Hokkaido University

This paper is posted at the eScholarship Repository, University of California.

<http://repositories.cdlib.org/ipnc/xvi/1051>

Copyright ©2009 by the authors.

Difference in Acquisition of Soil Organic Nitrogen between Pak-Choi and Tomato

Abstract

This study was to elucidate the factors contributing to organic nitrogen (N) acquisition in different plant species, pak-choi (*Brassica rapa* L. var. chinensis) and tomato (*Solanum lycopersicum* L.). Two hypotheses were proposed and examined: 1) higher protease activity in rhizosphere contributes to the ability to utilize organic N in soils; 2) the differences in ability to absorb amino acids are responsible for the different characteristics of organic N utilization. Seedlings were grown in soils mixed with vermiculite with different N sources (no N (-N), ammonium sulfate (AS), and cattle farmyard manure (CM)). Plant growth and N accumulation were the best in the AS treatment in both species. When comparing the growth and N accumulation between -N and CM treatments, those of pak-choi were significantly enhanced in the CM treatment, but no difference was found in tomato. In the rhizosphere soils, the highest protease activities were detected in the CM treatment irrespective of plant species. The correlation analysis between rhizospheric protease activity and plant total N accumulation in the -N and CM treatments showed that a significant positive correlation only in pak-choi. The determination of amino acid absorption rate in excised roots indicated that glycine was taken up at significantly higher rate in pak-choi than in tomato. Taken together, two possible factors for the acquisition of organic N was suggested from this study: protease activity in rhizosphere and ability to absorb amino acids.

Introduction

Effective utilization of organic matter is strongly desired to promote sustainable agriculture. Whereas it has been well known that growth responses to organic matter differ among plant species, the factors affecting these characteristics has not been elucidated. The most part of organic nitrogen (N) in organic matter is a proteaceous compound (Kumar et al., 2006). In soils, proteins are degraded by protease, and the released amino acids and peptides are mineralized to ammonium (NH_4) and nitrate (NO_3), which can be utilized by plant roots. Here, we propose two hypotheses accounting for the factors responsible for the differences in plant response to organic N. The first hypothesis is that higher protease activity in rhizosphere contributes to the ability to utilize organic N in soils. The rate-limiting step in degradation of organic nitrogen in soils is considered to be a protease-mediated process. Plant species with high ability to utilize organic N may make their rhizosphere environment more suitable for protease-producing microorganisms, or may secrete protease from roots, resulting in the enhancement of proteolysis in rhizosphere soils. The second hypothesis is that the differences in ability to absorb amino acids are responsible for the different characteristics of organic N utilization. Recently, it has been shown that many plant species can absorb amino acids as significant N-sources for their growth (Streeter et al., 2000). Plant species with high ability to absorb amino acids is considered to be superior characteristics when competing with rhizosphere microorganisms for N acquisition. In this study, plant utilization of organic N is comprehensively discussed by testing these hypotheses.

Materials and Methods

Pot experiment

Seedlings of pak-choi (*Brassica rapa* L. var. chinensis) and tomato (*Solanum lycopersicum* L.) were grown in pots in a greenhouse. Each pot contained 300 mL of vermiculite mixed with field soils, as sources of microbe inoculums (vermiculite : field soil = 4 : 1, v/v). Field soils were collected from Hokkaido Central Agricultural Experiment Station, Naganuma, Japan. For 10 years, 25000 kg of manure and 2000 kg of rice bran, and 1070 kg of fish meal $\text{ha}^{-1} \text{ year}^{-1}$ have been applied to the field. Different sources of N (no N (–N), ammonium sulfate (AS), and cattle farmyard manure (CM)) were applied to each soil for the N treatment (70 mg N pot^{-1} in AS and CM treatments).

After 28-days treatment, seedlings and soils (rhizosphere and non-rhizosphere soils) were sampled. Plant N concentrations were determined. Soil protease activities were determined according to Ladd and Bulter (1972), using benzyloxycarbonyl phenylalanyl leucine (ZFL) as a substrate.

Determination of amino acid uptake rate

Excised roots of pak-choi and tomato were incubated in 10 mL of 0.2 mM CaCl_2 solution (pH5.2) containing different sources of nitrogen (glycine (Gly), glutamic acid (Glu), NO_3^- and NH_4^+) for 2 hours at 25 °C or 4 °C. N concentration in the solution before and after the incubation was determined and the decrease in the N concentration was calculated. Active uptake rate for each N source was estimated by subtracting the rate at 4 °C (passive N uptake) from that at 25 °C.

Results and Discussion

Plant growth was the best in the AS treatment in both species (Fig. 1). In tomato, no significant difference in growth was found between the –N and CM treatments. In contrast, the growth of pak-choi was significantly enhanced in the CM treatment in comparison with the –N treatment. N accumulation in the plants showed the same tendency with plant growth (data not shown). These results suggest that pak-choi can utilize organic N in soils more efficiently than tomato.




Fig. 1. Dry weight of tomato and pak-choi seedlings after the treatment. –N, no N application; AS, ammonium sulfate; CM, cattle farmyard manure.

As described in introduction section, protease activity in rhizosphere soils is one of the key factors that contribute to organic N acquisition in plant. In the rhizosphere soils, the highest activities were detected in the CM treatment irrespective of plant species (data not shown), which suggested that the organic matter application increased the activities. Although rhizosphere effect (Badalucco et al., 1996) was observed for the protease activities when comparing the activities between rhizosphere and bulk soils, there was no difference in this effect between two species (data not shown). Then, the correlation between rhizospheric protease




Fig. 2. Relationship between N accumulation and rhizospheric protease activity in tomato and pak-choi seedlings. **, significant at $P < 0.01$

activity and plant total N accumulation was determined using individual replicates in both species grown with the –N and CM treatments (Fig. 2). A significant positive correlation was found only in pak-choi ($r^2=0.89$, $P<0.01$). These results suggest that the rhizospheric protease activity is a potential factor contributing to organic N acquisition in plant but other factors are likely to be involved in the mechanisms.

It is well known that plant roots compete with soil microbes for N acquisition in rhizosphere. It can be an advantage to take up amino acid N more efficiently to defeat rhizosphere microbes in N acquisition. Therefore, plant ability to absorb amino acid N may also be responsible for organic N acquisition in plant. As expected, Gly, which is one of the major free amino acid in soils (Streeter et al., 2000), was taken up at significantly higher rate in excised roots of pak-choi than of tomato (Table 1).

Table 1. Net uptake rate ($\mu\text{mol g}^{-1}$ root FW h^{-1}) of NO_3^- , NH_4^+ , Gly, and Glu in excised roots of tomato and pak-choi seedlings.

	NO_3^-	NH_4^+	Gly	Glu
Tomato	0.51 $\pm$ 0.16	1.37 $\pm$ 0.37	2.08 $\pm$ 0.31	2.15 $\pm$ 0.45
Pak-choi	0.12 $\pm$ 0.07	0.70 $\pm$ 0.42	5.55 $\pm$ 1.76	1.95 $\pm$ 1.02

In conclusion, this study strongly suggests that organic N acquisition in plant depends on at least the following two factors; protease activity in rhizosphere and ability to absorb amino acids. It would be interesting to characterize the microbiota in rhizosphere from the point of view of protease production and/or competition in amino acid acquisition.

References

- Badalucco L, Kuikman PJ, Nannipieri P, Protease and deaminase activities in wheat rhizosphere and their relation to bacterial and protozoan populations. *Biology and Fertility of Soils*. 1996; 23: 99-104.
- Kumar K, Rosen CJ, Russelle MP, Enhanced protease inhibitor expression in plant residues slows nitrogen mineralization. *Agronomy Journal*. 2006; 98: 514-521.
- Ladd JN, Bulter JHA, Short-term assays of soil proteolytic enzyme activities using proteins and dipeptide derivatives as substrates. *Soil Biology and Biochemistry*. 1972; 4: 19-30.
- Streeter TC, Bol R, Bardgett RD, Amino acids as a nitrogen source in temperate upland grasslands: the use of dual labelled (^{13}C , ^{15}N) glycine to test for direct uptake by dominant grasses. *Rapid Communications in Mass Spectrometry*. 2000; 14: 1351-1355.