

NENT-D-16-00260R1

Disjunct occurrence of *Trichadenotecnum s.str.* in southeastern Brazil  
(Psocodea: "Psocoptera": Psocidae), with description of a new species

K Yoshizawa 1, M Souza-Silva 2, RL Ferreira 2

1 Systematic Entomology, School of Agriculture, Hokkaido Univ, Sapporo, Japan

2 Centro de Estudos em Biologia Subterrânea, Setor de Zoologia Geral, Departamento de  
Biologia, Universidade Federal de Lavras, Lavras, Brazil

**Keywords**

psocid, new species, disjunct distribution, biogeography, South America

**Correspondence**

K Yoshizawa, Systematic Entomology, School of Agriculture, Hokkaido Univ, Sapporo  
060-8589, Japan; psocid@res.agr.hokudai.ac.jp

**Abstract**

A new barklouse species belonging to the *Trichadenotecnum s.str.* clade (Psocodea: "Psocoptera": Psocidae) was described from southeastern Brazil (Minas Gerais state). This locality is highly isolated (about 3400 km) from the closest known distributional range of this clade. It also represents the southern-most distributional record of *Trichadenotecnum s.str.* Phylogenetic placement of the species and biogeography of Neotropical *Trichadenotecnum* were briefly discussed. *Trichadenotecnum punctipenne* New, 1972 described from Brazil was synonymized with *T. pardus* Badonnel, 1955.

## Introduction

The genus *Trichadenotecnum* is one of the most species-rich and widely distributed barklouse genera (Psocodea: "Psocoptera"). It occurs in South America but, in comparison to the high diversification of the genus in Southeast to eastern Asia (ca. 150 species: Yoshizawa *et al*, 2014; Yoshizawa & Lienhard, 2015a, b) or in Central America (over 35 species: Yoshizawa *et al*, 2008), species richness of *Trichadenotecnum* in South America is quite low (10 species: Yoshizawa *et al*, 2008; Yoshizawa & García Aldrete, 2010). Recent morphological (Yoshizawa *et al*, 2008) and molecular (Yoshizawa *et al*, 2016) phylogenetic analyses showed that most of the South American *Trichadenotecnum* species belong to a single small clade (Neotropical Clade in Fig 1) which is the sister to all the rest of members of the genus (Yoshizawa *et al*, 2008, 2016). Within the major clade of the genus, a conspicuous subclade, *Trichadenotecnum s.str.*, has long been recognized (Fig 1), which is distributed throughout the Oriental and Holarctic Regions and throughout the northern-most part of the Neotropics (Figs 1–2). *Trichadenotecnum s.str.* has its origin in the Old World (probably Oriental), and the occurrence of the species of this clade in the New World is considered to be a result of transcontinental dispersals from the Old World (Yoshizawa *et al*, 2008). Only two species of *Trichadenotecnum s.str.* are known to occur in South America: *T. decui* Badonnel, 1987 from Venezuela and *T. simile* Mockford, 1996, which was originally described as the male of *T. decui* by Mockford (1991) but was later recognized as an independent species, from northern Brazil. Both species have their close relatives in Central America (North, Central, South American Clade in Fig 1: Yoshizawa *et al*, 2008). Extremely low diversification of *Trichadenotecnum s.str.* in South America and its restriction to the northern-most region of the continent have been considered to reflect their recent dispersal to South America, after formation of the Central and South American connection in the Pliocene (ca. 3 Ma) (Yoshizawa *et al*, 2008).

In this paper, we report the occurrence of a species belonging to *Trichadenotecnum s.str.* from southeastern Brazil (Lavras, in southern Minas Gerais state), which is here described as *Trichadenotecnum ufla* n. sp. (Fig 3A). This locality is over 3400 km away from the known localities of the *Trichadenotecnum s.str.* species in South America (Venezuela and Roraima state in northern Brazil) (Fig 2). It also represents the southern-most distributional record of the clade.

## Material and Methods

The specimens studied were collected by beating and were freshly killed and stored in 95% ethanol. Examination was made by using a stereoscopic microscope (SZX16: Olympus, Japan) or a compound microscope (Axiophot: Carl Zeiss Microscopy, Germany). The photographs were taken by using a digital camera (OM-D E-M5: Olympus, Japan) attached to SZX 16 (habitus) or Axiophot (forewing). Terminology followed Yoshizawa *et al* (2014).

## Taxonomy

### *Trichadenotecnum ufla* Yoshizawa, n. sp. (Figs 3–4)

#### *Description*

*Female* (Fig 3A). Head. White in ground color; vertical markings blackish brown, each marking separated from neighboring marks; orbital markings brown, posterior spots paler; brown marking between vertical and orbital markings; coronal suture white; epicranial suture bordered with brown band, outer part broader; frons with two pair of dark brown markings and medially with small spot; eyes gray, IO/D = 2.1; ocelli white, ocellar field black; gena blackish brown, medially with pale portion; postclypeus with rows of dark brown spots dorsally, spots fused and forming brown marking ventromedially; anteclypeus blackish brown. Scape and pedicel dark brown; flagellum pale brown. Mouthparts blackish brown.

Thorax. Prothorax pale brown. Mesonotum white in ground color, with posterior part of lateral lobe brown; median lobe mostly brown, sutures blackish brown. Metanotum mostly white, brown medially. Meso- and metapleuron blackish brown except for white membranous region.

Legs. Fore- and midlegs mostly white; coxa of midleg blackish; femora with two blackish stripes; tibiae with blackish stripe near distal tip. Hindleg blackish except for white tibia with blackish stripe near distal tip. Tarsi of all legs blackish.

Forewing (Fig 3B). Spots in cell a1 well developed, basal one paler. Opposing spots in cell r apparent, about equal in size, clearly separated. Basal band discontinuous around M–Cu

1 fork, with broad oval pale region in cell cu. Median spots well developed. Distal band well  
2 developed. Spot on roof of cell m3 broad. Submarginal spots well developed, especially in  
3 cell r5 expanding distally to marginal cloud. Marginal cloud dark and well developed, with  
4 pale region at wing margin of cells r5 and m1 and middle of cell m3.  
5

6  
7  
8 Abdomen pale yellow with small brown irregular markings dorsally, brown ventrally.  
9

10 Genitalia (Fig 4). Subgenital plate (Fig 4A) with pair of pointed and basally serrated  
11 processes posteriorly; egg guide weakly sclerotized ventrally, widely sclerotized dorsally,  
12 posterior margin rounded, narrow and parallel sided in distal half, then gradually broadened  
13 to base; body broadly sclerotized with narrow membranous region medially, posterolaterally  
14 with wrinkled region just anterior to long processes, and with extended sclerotization  
15 anteromedially. Gonapophyses (Fig 4B). Ventral valve long; dorsal valve with long distal  
16 process; external valve with broad posterior lobe, internal lobe short. Spermapore plate  
17 sclerotized and pigmented as in Fig 4C.  
18  
19  
20  
21  
22  
23  
24  
25

26  
27 *Measurements.* Body length 2.1–2.2 mm; forewing length 2.8–2.9 mm; hindwing length 2.0–  
28 2.1 mm.  
29  
30

31  
32 *Type material.* Holotype: female. S21.223747°, W44.968633°, Campus of the Universidade  
33 Federal de Lavras (UFLA), 16.iii.2016, K. Yoshizawa (deposited in ISLA-UFLA) (five  
34 sequences, *16S rRNA*, *12S rRNA*, *COI*, *18S rRNA* and *Histone 3* genes, of the holotype are  
35 deposited in GenBank, under the accession numbers LC185091–5). *Paratypes.* 2 females,  
36 same locality as for holotype, 1 collected on 16.iii and 1 collected on 17.iii.2016 (deposited in  
37 ISLA-UFLA and Hokkaido University)  
38  
39  
40  
41  
42  
43  
44

45 *Etymology.* The specific epithet, an invariable combination of letters, is the abbreviation of  
46 the type locality, Universidade Federal de Lavras.  
47  
48  
49

50  
51 *Habitat.* *Trichadenotecnum ufla* was collected in a secondary forest (Fig 5A: of about 7.56  
52 ha) predominantly formed by trees of *Anadenanthera peregrina*, with an understory of  
53 diversified vegetation (at least 25 families and 66 plant species) (Fig 5B). This forest resulted  
54 from a shading experiment of coffee trees (*Coffea arabica*) started in 1943, currently in a  
55 permanent preservation area of UFLA (Nascimento *et al*, 2009). This forest is surrounded by  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 *Eucalyptus* plantations, public roads, forest nurseries, coffee plantations, and buildings of the  
2 Coffee Research Centre (Fig 5A).

3  
4  
5 The local climate has rainy and hot dry summers and cold winters, with average annual  
6 rainfall above 1400 mm (ranging from 15 to 3000mm) and average monthly temperatures  
7 from 15 to 23°C (Dantas *et al*, 2007). The rains occur between November and February  
8 (Dantas *et al*, 2007; INMET 2016).  
9

10  
11  
12  
13  
14 *Remarks.* This species belongs to *Trichadenotecnum s.str.* and resembles the species of the  
15 *alinguum* group distributed in the Oriental Region (not presented in Fig 1: see Yoshizawa &  
16 Lienhard, 2015b and Yoshizawa *et al*, 2016 for the morphology and placement of the species  
17 group) in having long and pointed processes on the subgenital plate. However, preliminary  
18 molecular phylogenetic analysis (Yoshizawa *et al*, unpublished) shows that this species is  
19 closely related to the New World species of the genus (see below). Among the New World  
20 species of *Trichadenotecnum*, nothing similar is known.  
21

22  
23  
24  
25  
26  
27 *T. simile* Mockford, 1996 is known to occur in Brazil (Roraima, Rio Uraricoera, Ilha de  
28 Maraca), the female of this species has not been recorded to date. However, the forewing  
29 markings of *T. ufla* (Fig 3B) differ significantly from those of *T. simile* in the following points  
30 (see Mockford, 1991: fig. 287): opposing spots in cell r are about equal in size in *T. ufla*  
31 whereas the posterior spot is much smaller in *T. simile*; the submarginal spot in cell m3 is  
32 distinct in *T. ufla* whereas it is fused to the marginal cloud and m3 roof spot so that is obscure  
33 in *T. simile*; marginal cloud is discontinuous in cell m2 in *T. ufla* whereas it is continuous in  
34 *T. simile*. The forewing marking condition is quite stable at least among the three individuals  
35 of *T. ufla* examined here. Forewing length of *T. simile* (2.2 mm) is much smaller than that of  
36 *T. ufla*, although such variation can sometimes be seen even within the same species.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

## 47 **Discussion**

48  
49  
50  
51 Of two known species of the South American *Trichadenotecnum s.str.* clade, *T. decui* is  
52 known from both sexes whereas *T. simile* is known only from a male. Therefore, there is a  
53 possibility that the present species, known so far only from females, may be conspecific with  
54 *T. simile*. However, from the differences in the forewing markings (see *Remarks* above) and  
55 great geographical isolation with climate differences (Fig 2), we concluded that these  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

specimens represent different species. Description of a *Trichadenotecnum* species based only on the females may be criticized, but DNA sequences (five genes, including a partial mitochondrial *COI*, the commonly used marker for DNA barcoding) of the holotype are available from the public database (see above for GenBank accession number) so that future identification of the conspecific male is warranted.

The subgenital plate of *T. ufla* is characteristic in having a pair of long and narrow processes, unknown in any other New World *Trichadenotecnum* species. Similar structures, however, can be seen in some species of the Oriental *Trichadenotecnum* (the *alinguum* group and a couple of Nepalese species of the *spiniserrulum* group: Yoshizawa *et al*, 2007; Yoshizawa & Lienhard, 2015b). Therefore, at least three independent origins of such processes can be assumed. The function of the processes is unknown to date.

Preliminary molecular analyses (Yoshizawa *et al*, unpublished) placed this species to the basal branch of the *slossonae* group (Fig 1), in which *T. simile* is classified (Yoshizawa *et al*, 2008). The morphology-based phylogenetic analysis supported sister group relationship between the *slossonae* and *decui* groups (Fig 1), but no species of the *decui* group was sampled in the molecular analyses. Therefore, as to the species-group placement of *T. ufla*, these two species groups are both probable. Here, we tentatively assign this species to the *slossonae* group because of the similarity of the spermapore plate structure of this species with that of *T. slossonae* Banks, 1903. The subgenital plate of *T. slossonae* possesses weakly projecting parts on the posterior margin, which probably represent the ancestral condition of the well-developed processes on the subgenital plate of *T. ufla*.

Although *Trichadenotecnum s.str.* is quite diversified in Central America, with a total of 35 species classified into seven species groups (North, Central, South American Clade in Fig 1), all three South American species of *Trichadenotecnum s.str.* belong to a single clade (the *slossonae* + *decui* clade). This may reflect the biogeographical history of *Trichadenotecnum s.str.* in Central and South America, but further phylogenetic and biogeographic analyses are needed to uncover when *Trichadenotecnum s.str.* colonized South America, why South American species are restricted to a single clade, and how they expand their distributional range in the continent. The widely disjunct distribution of *Trichadenotecnum s.str.* in South America is interesting because it potentially reflects the Quaternary refugia (Brown *et al*, 1974; Dietz, 1994). However, it may be just due to poor faunal investigation of Psocoptera in South America. Further investigation of the *Trichadenotecnum* fauna in South America,

1 especially for the high altitude regions rather than the Amazonian area, is highly desirable,  
2 because occurrence of *Trichadenotecnum s.str.* in South America seems to be restricted to  
3 mountainous regions (Fig 2).  
4  
5

6  
7 The present discovery also highlights the importance of preserved forested areas in  
8 urban centers (Fig 5A). Unfortunately, during the expansion of many cities, most forested  
9 areas are removed, and many habitats (especially microhabitats) of some species can be  
10 destroyed. Most forested areas in cities (usually parks or squares) are created to promote  
11 recreation for the population, but without the concern of maintaining original habitats for  
12 different species. Especially in the Neotropics, with well-known mega diversity, it is crucial  
13 to keep forested areas in cities, since they can certainly embrace the main habitat of many  
14 species, particularly endemic invertebrates. In the case of the Universidade Federal de  
15 Lavras, the preserved areas have already revealed a new species, the amphipod *Hyalrella*  
16 *minensis* (Bastos-Pereira & Bueno, 2013).  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

27 **Acknowledgments** We thank Luiz Felipe Moretti Iniesta and Yoshitaka Kamimura for  
28 support in the field, Mateus Lima da Silva for providing the satellite image of UFLA, and  
29 Izumi Yao for support of DNA experiments. This study was partly supported by JSPS  
30 research grants (24570093 and 15H04409) to KY and National Council of Technological and  
31 Scientific Development (CNPq) (No. 304682/2014-4) to RLF.  
32  
33  
34  
35  
36  
37

## 38 **References**

39  
40  
41

- 42 Badonnel A (1955) Psocoptères de l'Angola. Publicações culturais Companhia Diamantes  
43 Angola 26:1–267  
44  
45 Badonnel A (1987) Psocoptères de Venezuela et de la République Argentine. In: Decu V,  
46 Orghidan T, Bordon C, Linares O, Urbani F, Tronchoni J & Bosque C. Fauna hipogea y  
47 hemiedáfica de Venezuela y de otros países de America del Sur. Editura Academiei  
48 Republicii Socialiste Romania, Bucharest, 173–182  
49  
50 Banks N (1903) Some new Neuropteroid insects. J NY Entomol Soc 11:236–243  
51  
52 Bastos-Pereira R, Bueno, AAP (2013) A new species of freshwater amphipod  
53 (Dogielinotidae, *Hyalrella*) from Southeastern Brazil. Nauplius 21:79-87  
54  
55 Brown KS, Sheppard PM, Turner JRG (1974) Quaternary refugia in tropical America:  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- evidence from race formation in *Heliconius* butterflies. Proc R Soc Lond B 187:369–378
- Dantas AAA, Carvalho LG, Ferreira E (2007) Climatic classification and tendencies in Lavras region, MG. Ciência agrotecnologia 6:1862-1866
- Dietz RE (1994) Systematics and biology of the genus *Macrocneme* Hübner (Lepidoptera: Ctenuchidae). Univ California Publ Entomol 113:1–121
- INMET (2016) Instituto Nacional de Meteorologia. [www.inmet.gov.br](http://www.inmet.gov.br)
- Mockford EL (1991) New species and records of Psocoptera (Insecta) from Roraima state, Brazil. Acta Amazonica 21:211–318
- Mockford EL (1993) North American Psocoptera (Insecta). Flora Fauna Handbook 10. Sandhill Crane Press, Florida, p 455
- Mockford EL (1996) New species and records of Psocoptera from northern Venezuela. Tropical Zool, Special Issue 2:1–98
- Nascimento GA, Pifano DS, Lima MP, Calegário N (2009) Floristic aspects and diversity of regenerated arboreal species under a stand of *Anadenanthera peregrina* Speg. Cerne, Lavras 2(15):187-195
- New TR (1972) Some Brazilian Psocoptera from bird nests. Entomologist 105:153–160
- Yoshizawa K, Lienhard C, Thapa VK (2007) Systematic study of the genus *Trichadenotecnum* in Nepal (Psocodea: 'Psocoptera': Psocidae). Insecta matsumurana NS 63:1–33
- Yoshizawa K, García Aldrete AN, Mockford EL (2008) Systematics and biogeography of the New World species of *Trichadenotecnum* Enderlein (Insecta: Psocodea: 'Psocoptera': Psocidae). Zool J Linn Soc 153:651–723
- Yoshizawa K, García Aldrete AN (2010) Description of a new species of the *Trichadenotecnum roesleri* species group from Brazil (Insecta: Psocodea: 'Psocoptera': Psocidae) and its phylogenetic implications. Stud Neotr Fauna Env 45: 61-66
- Yoshizawa K, Lienhard C, Idris AB (2014) *Trichadenotecnum* species from Peninsular Malaysia and Singapore (Insecta: Psocodea: 'Psocoptera': Psocidae). Zootaxa 3835:469–500
- Yoshizawa K, Lienhard C (2015a) Synonymy of *Cryptopsocus* Li with *Trichadenotecnum* Enderlein (Insecta: Psocodea: 'Psocoptera': Psocidae) and description of three new species. Zootaxa 3957:480–488



1 Yoshizawa K, Lienhard C (2015b) The *alinguum* group, a new species group of the genus  
2  
3 *Trichadenotecnum*, with description of two new species from Thailand (Psocodea:  
4  
5 'Psocoptera': Psocidae). Insecta matsumurana NS 71:1789–188  
6

7 Yoshizawa K, Yao I, Lienhard C (2016) Molecular phylogeny reveals genital convergences  
8  
9 and reversals in the barklouse genus *Trichadenotecnum* (Insecta: Psocodea:  
10  
11 'Psocoptera': Psocidae). Mol Phylog Evol 94:358–364  
12  
13  
14  
15

## 16 **Appendix. Identity of *Trichadenotecnum punctipenne* New, 1972**

17  
18 (by K. Yoshizawa)  
19  
20

21  
22 In addition to the species mentioned in the main text of this paper, one more species, *T.*  
23 *punctipenne* is also known to occur in Brazil (Pernambuco state). Judging from a unique  
24  
25 structure on the subgenital plate (lining sclerotisations), this species is apparently close to *T.*  
26  
27 *pardus* Badonnel, 1955 and *T. pardidum* Thornton, 1961. The former is parthenogenetic and  
28  
29 is widely distributed worldwide including Brazil (Mockford, 1993; Yoshizawa *et al*, 2008).  
30  
31 This wide distribution of the parthenogenetic species is considered to be artificial, introduced  
32  
33 from its native area in the Oriental Region, where the sister species of *T. pardus* (i.e., bisexual  
34  
35 *T. pardidum*) is distributed.

36  
37 Judging from the original description, I could not find any difference between *T.*  
38  
39 *punctipenne* and *T. pardus*. All close relatives of *T. punctipenne*, i.e., species of the  
40  
41 *spiniserrulum* group (Fig 1), are unknown from the New World. Very isolated occurrence of a  
42  
43 native species of the *spiniserrulum* group (distributed in the Oriental and Afrotropical  
44  
45 Regions: Fig 1) in Brazil is less likely. Therefore, I conclude here that *T. punctipenne* should  
46  
47 be treated as a **new junior synonym** of *T. pardus*.

48  
49 As mentioned above, *T. pardus* is likely to be artificially introduced to South America.  
50  
51 Therefore, its occurrence in central Brazil does not affect the biogeographical discussion  
52  
53 made in the main text of this paper.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Figure captions

Fig 1 Phylogenetic relationships among representative species groups of *Trichadenotecnum* (Yoshizawa *et al*, 2008). Note that some Old World species groups (e.g., the *alinguum* group) are not included in the tree (see Yoshizawa *et al*, 2016 for the placement of such groups). Distributional range of each species group (except for artificially introduced ones) is indicated after the group name. AF: Africa; CA: Central America; EP: eastern Palearctic; HA: Holarctic; NA: North America; OR: Oriental; PA: Palearctic; SA: South America; WP: western Palearctic.

Fig 2 Distribution map of the Neotropical species of *Trichadenotecnum s.str.*

Fig 3 *Trichadenotecnum ufla* n. sp. *A* Habitus of paratype female; *B* forewing of the holotype female.

Fig 4 Female genitalia of *Trichadenotecnum ufla* n. sp., ventral view. *A* Subgenital plate; *B* gonapophyses; *C* Spermapore plate.

Fig 5 The type locality of *Trichadenotecnum ufla* n. sp. *A* Satellite view: red broken line indicates the forest area where the species was collected; *B* environment within the forest.










Figure 4




Figure 5

[Click here to download colour figure Fig.5.tif](#)

